

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
24 December 2003 (24.12.2003)

PCT

(10) International Publication Number  
WO 03/106402 A1

not Sk

(51) International Patent Classification<sup>7</sup>: C07C 235/34,  
C07D 209/20, 213/55, 333/24, A61K 31/197, 31/405,  
A61P 13/00

(21) International Application Number: PCT/EP03/05705

(22) International Filing Date: 30 May 2003 (30.05.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0213488.0 12 June 2002 (12.06.2002) GB(71) Applicant (for all designated States except US): BAYER  
AKTIENGESELLSCHAFT [DE/DE]; 51368 Lev-  
erkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SHIMAZAKI,  
Makoto [JP/JP]; 5-5-3-1, Kunimidai, Kizu-cho, So-  
raku-gun, Kyoto 619-0216 (JP). SAKURAI, Osamu  
[JP/JP]; 11-15, Koaza-Hiroshiki, Aza-Oyamazaki,  
Oyamazaki-cho, Otokuni-gun, Kyoto 618-0071 (JP).  
MURATA, Toshiki [JP/JP]; 1-3-501, Nakano-cho,  
Ikoma-shi, Nara 630-0267 (JP). URBHANS, Klaus  
[DE/JP]; 6-3-1-301, Kusugaoka-cho, Nada-ku, Kobe,  
Hyogo 657-0024 (JP). YAMAMOTO, Noriyuki [JP/JP];  
6-6-1-3-404, Jingu, Nara-shi, Nara 631-0804 (JP).  
YOSHIKAWA, Satoru [JP/JP]; 4-10-6, Shibatsuji-cho,  
Nara-shi, Nara 630-8114 (JP). UMEDA, Masaomi  
[JP/JP]; 2-840-156, Sahodai, Nara-shi, Nara 630-8105  
(JP). TAJIMI, Masaomi [JP/JP]; 1-8-17, Sakuragaoka,  
Seika-cho, Soraku-gun, Kyoto 619-0232 (JP). MASUDA,  
Tsutomu [JP/JP]; 3-15-6-6A, Jingu, Nara-shi, Nara  
631-0804 (JP). SHINTANI, Takuya [JP/JP]; 4-24-1,  
Seikadai, Seika-cho, Soraku-gun, Kyoto 619-0238 (JP).  
SHIMIZU, Haruka [JP/JP]; 88, B-202, Aono-cho,  
Nara-shi, Nara 631-0841 (JP).(74) Common Representative: BAYER AKTIENGE-  
SELLSCHAFT; 51368 Leverkusen (DE).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU;  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,  
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,  
UZ, VC, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,  
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU,  
SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG,  
UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR),  
OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG)

## Published:

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendmentsFor two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: 2-NAPHTHAMIDE DERIVATIVES

RCK 29

(57) Abstract: The present invention relates to 2-naphthamides, which are useful as an active ingredient of pharmaceutical prepa-  
rations. The 2-naphthamides of the present invention have IP receptor antagonistic activity, and can be used for the prophylaxis and  
treatment of diseases associated with IP receptor activity. Such diseases include urological diseases or disorder as follows: bladder  
outlet obstruction, overactive bladder, urinary incontinence, detrusor hyper-reflexia, detrusor instability, reduced bladder capacity,  
frequency of micturition, urge incontinence, stress incontinence, bladder hyperreactivity, benign prostatic hypertrophy (BPH), pro-  
statitis, urinary frequency, nocturia, urinary urgency, pelvic hypersensitivity, urethritis, pelvic pain syndrome, prostatodynia, cystitis,  
or idiopathic bladder hypersensitivity. The compounds of the present invention are also useful for treatment of pain including, but not  
limited to inflammatory pain, neuropathic pain, acute pain, chronic pain, dental pain, premenstrual pain, visceral pain, headaches,  
and the like; hypotension; hemophilia and hemorrhage; and inflammation, since the disease is also alleviated by treatment with an  
IP receptor antagonist.

BEST AVAILABLE COPY

WO 03/106402 A1



## 2-Naphthamide derivatives

### Detailed Description of Invention

#### Technical Field

The present invention relates to a 2-naphthamides which are useful as an active ingredient of pharmaceutical preparations. The 2-naphthamides of the present invention have an IP receptor antagonistic activity, and can be used for the prophylaxis and treatment of diseases associated with IP receptor activity.

More specifically the 2-naphthamide derivatives of the present invention are useful for treatment and prophylaxis of urological diseases or disorders.

The compounds of the present invention are also useful for treatment of pain; hypotension; hemophilia and hemorrhage; inflammation; respiratory states from allergies or asthma, since the disease also is alleviated by treatment with an IP receptor antagonist.

#### BACKGROUND ART

Prostaglandins (or prostanoids, PGs) are a group of bioactive lipid mediators generated from membrane phospholipids. They are formed from 20-carbon essential fatty acids containing 3, 4, or 5 double bonds, and carry a cyclopentane ring. They are divided into 6 main classes (D, E, F, G, H or I) by the cyclopentane ring structure. The main classes are further subdivided by subscripts 1, 2, or 3, reflecting their fatty acid precursors. PGI<sub>2</sub> is a member of prostanoids, and it has a double ring structure and is derived from arachidonic acid. The receptor for PGI<sub>2</sub> is a seven transmembrane G-protein coupled receptor, called IP receptor. IP receptor couples at least to G<sub>s</sub>-type G-protein, and activates adenylate cyclase and phospholipase C. The

expression of IP receptor is demonstrated in aorta, coronary/pulmonary/cerebral arteries, platelets, lung, and dorsal root ganglions in addition to several other tissues.

5 One of the well-known actions of PGI<sub>2</sub> on blood vessels is to cause vasodilation and hypotension. Especially in septic shock, PGI<sub>2</sub> is produced and participate in the induction of systemic hypotension (G.D. Bottoms et al, Am J Vet Res 1982, 43(6), 999-1002). Therefore, IP receptor antagonists may prevent hypotension associated with septic shock.

10 Another well-known action of PGI<sub>2</sub> on platelets is to suppress aggregation. In the IP receptor knock out mice, FeCl<sub>3</sub>-induced thrombosis formation was enhanced in comparison with that in wild type mice (T. Murata et al, Nature 1997, 388, 678-682.), confirming the involvement of IP receptor in the platelet inhibition. Therefore, IP receptor antagonists may enhance the platelet activation and suppress excessive  
15 bleeding such as, but not limited to, hemophilia and hemorrhage.

PGI<sub>2</sub> also participate in the inflammation. In the inflamed tissue, various inflammatory mediators, including prostaglandins, are produced. PGI<sub>2</sub> is also generated and induces vasodilation to increase blood flow. This enhances vascular permeability,  
20 edema formation and leukocyte inflammation in the inflamed region (T. Murata et al, Nature 1997, 388, 678- 682.). Therefore, IP receptor antagonists may be efficacious for the treatment of inflammation.

PGI<sub>2</sub> may be involved in the pathogenesis of respiratory allergy or asthma. It is  
25 spontaneously generated and the major prostaglandin in human lung, and the appropriate antigen challenge increases PGI<sub>2</sub> production (E.S. Schulman et al, J Appl Physiol 1982, 53(3), 589-595.). Therefore, IP receptor antagonists may have a utility for the treatment of those respiratory diseases.

30 In addition, an important role of IP receptor in the induction of hyperalgesia has been clearly shown by IP receptor knockout mice (T. Murata et al., Nature 1997, 388,


678-682). Injection of acetic acid into the peritoneal cavity induced production of PGI<sub>2</sub>. This PGI<sub>2</sub> is considered to bind to IP receptor on sensory neurons. As IP receptor couples to the activation of both adenylate cyclase and phospholipase C, cAMP-dependent protein kinase (PKA) and protein kinase C (PKC) are activated. PKA and PKC are known to modulate ion channels on sensory neurons such as VR1, P2X<sub>3</sub>, and TTX-R. As a result, PGI<sub>2</sub> sensitizes sensory neurons to enhance the release of neurotransmitters. Hence, acetic acid injection induces nociceptive response (writhing) in mice. This acetic acid-induced writhing was greatly reduced in PGI<sub>2</sub> receptor-null mice as the same level as indomethacin-treated wild type mice. Several other in vivo hyperalgesia studies in rodents and in vitro studies further support that PGI<sub>2</sub> plays a major role in the induction of hyperalgesia and that PGI<sub>2</sub> acts as important modulator of sensory neurons (K. Bley et al, Trends in Pharmacological Sciences 1998, 19(4), 141-147.). Therefore, IP receptor antagonists may be useful for the treatment of pain.

Sensory neurons play very important roles not only in the pain sensation but also in the sensation of bladder distension. In normal subjects, A-delta sensory fibers are considered to play a major role to sense the bladder distention. However, in disease conditions of overactive bladder by, but not limited to, spinal cord injury, cystitis, Parkinson's disease, multiple sclerosis, previous cerebrovascular accident, and bladder outlet obstruction (BOO) caused by benign prostate hyperplasia (BPH), the sensitivity of C-fiber sensory neurons is upregulated and they contribute to the induction of the lower urinary tract symptoms. Treatment of overactive bladder patients with intravesical injection of capsaicin or its potent analog, resiniferatoxin, both of which desensitize VR1-positive C-fiber afferent neurons innervating the bladder, has been shown to be efficacious in several clinical trials (C. Silva et al, Eur Urol. 2000, 38(4), 444-452.). Therefore, C-fiber sensory neurons play an important role in the pathology of overactive bladder. PGI<sub>2</sub> is generated locally in the bladder and it is the major prostaglandin released from the human bladder. In a rabbit BOO model, a stable metabolite of PGI<sub>2</sub> was reported to be increased in BOO bladder (JM. Masick et al, Prostaglandins Other Lipid Mediat. 2001, 66(3), 211-219.).

Hence, PGI<sub>2</sub> from disease bladder sensitizes C-fiber sensory neurons, and as a result, it may induce symptoms of overactive bladder. Therefore, IP receptor antagonists are expected to be useful in the treatment of overactive bladder and related urinary disorders.


5

WO 00/31045 discloses anti-thrombotics agents represented by the general formula:




10

WO 98/44797 discloses integrin antagonists and farnesyl protein transferase inhibitors represented by the general formula:



15

EP-A-220 118 discloses pharmaceutical composition intended for the treatment of dermatological, respiratory and ophthalmological conditions represented by the general formula:



However, none of the references and other reference discloses 2-naphthamide derivatives having IP receptor antagonistic activity.

5

The development of a compound, which has effective IP receptor antagonistic activity and can be used for the prophylaxis and treatment of diseases alleviated by treatment with an IP receptor antagonist, has been desired.

10

#### Summary of the invention


15

As the result of extensive studies on chemical modification of 2-naphthamide derivatives, the present inventors have found that the compounds of the structure related to the present invention have unexpectedly excellent IP receptor and/or antagonistic activity. The present invention has been accomplished based on these findings.

This invention is to provide a novel 2-naphthamide derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof:

20

- (1) A 2-naphthamide derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof:



(I)

wherein

m and n independently represent an integer from 0 to 2;

5         $-R^1$  represents  $-O-R^{10}-OR^{11}$ ,  $-OR^{11}$ ,  $-SR^{11}$ ,  $-S(O)R^{11}$ ,  $-S(O)_2R^{11}$ ,  $-NR^{12}R^{13}$ ,  
or  $-CHR^{14}R^{15}$ ,

wherein

10         $-R^{10}$  represents  $(C_{1-6})$  alkylene;

15         $R^{11}$  represents aryl,  $(C_{2-6})$  alkenyl optionally substituted by aryl or  
heteroaryl,  $(C_{2-6})$  alkynyl optionally substituted by aryl or  
heteroaryl, or  $(C_{1-6})$  alkyl optionally substituted by  $(C_{3-8})$ -  
cycloalkyl, aryl or heterocycle comprising 4-9 carbons and at  
least one N, O, or S as a heteroatom,

wherein

20        said  $(C_{3-8})$  cycloalkyl, aryl and heterocycle optionally have one  
or two substituents selected from the group consisting of  
halogen, hydroxy, nitro,  $(C_{1-6})$  alkyl optionally substituted by  
mono-, di-, or tri halogen, and  $(C_{1-6})$  alkoxy optionally  
substituted by  $(C_{3-8})$  cycloalkyl, or mono-, di-, or tri halogen;

25         $R^{12}$  and  $R^{13}$  independently represent hydrogen,  $(C_{2-6})$  alkenyl  
optionally substituted by aryl or heteroaryl,  $(C_{2-6})$  alkynyl  
optionally substituted by aryl or heteroaryl, or  $(C_{1-6})$  alkyl  
optionally substituted by aryl or heteroaryl,

30        or



$R^{12}$  and  $R^{13}$  form, together with the nitrogen atom, a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

$R^{14}$  and  $R^{15}$  independently represent hydrogen,  $(C_{2-6})$ alkenyl optionally substituted by aryl or heteroaryl,  $(C_{2-6})$ alkynyl optionally substituted by aryl or heteroaryl,  $(C_{1-6})$  alkyl optionally substituted by aryl or heteroaryl, or  $(C_{1-6})$  alkoxy optionally substituted by aryl or heteroaryl,

or

$R^{14}$  and  $R^{15}$  form, together with the CH, a  $(C_{3-8})$ cycloalkyl optionally interrupted by NH, or O, or a phenyl optionally substituted by hydroxy, halogen or  $(C_{1-6})$  alkyl;

$R^2$  represents hydrogen, hydroxy, cyano,  $(C_{1-6})$  alkoxy,  $(C_{2-6})$ alkenyl,  $(C_{2-6})$ alkynyl,  $(C_{3-7})$ cycloalkyl, or  $(C_{1-6})$  alkyl optionally having one or two substituents selected from the group consisting of hydroxy, amino,  $(C_{1-6})$ alkylamino, aryl, and heteroaryl comprising 4-10 carbons and at least one N, O, or S as a heteroatom,

wherein

said aryl and heteroaryl optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, amino,  $N((C_{1-6})$  alkyl sulfonyl)amino, morpholino, phenyl, pyridyl,  $(C_{1-6})$  alkoxy optionally substituted by mono-, di-, or tri halogen, and  $(C_{1-6})$  alkyl optionally substituted by mono-, di-, or tri halogen; and

$R^3$  represents hydrogen, or  $(C_{1-6})$  alkyl.

The compounds of the present invention show excellent IP receptor antagonistic activity. They are, therefore, suitable for the production of medicament or medical composition, which may be useful to treat IP receptor related diseases.

5 More specifically, since the 2-naphthamide derivatives of the present invention antagonize IP receptor, they are useful for treatment and prophylaxis of urological diseases or disorder.

10 Such diseases or disorders include bladder outlet obstruction, overactive bladder, urinary incontinence, detrusor hyper-reflexia, detrusor instability, reduced bladder capacity, frequency of micturition, urge incontinence, stress incontinence, bladder hyperreactivity, benign prostatic hypertrophy (BPH), prostatitis, urinary frequency, nocturia, urinary urgency, pelvic hypersensitivity, urethritis, pelvic pain syndrome, prostatodynia, cystitis, or idiopathic bladder hypersensitivity.


15

The compounds of the present invention are also useful for treatment of pain including, but not limited to inflammatory pain, neuropathic pain, acute pain, chronic pain, dental pain, premenstrual pain, visceral pain, headaches, and the like; hypotension; hemophilia and hemorrhage; inflammation; respiratory states from  
20 allergies or asthma, since the disease is also alleviated by treatment with an IP receptor antagonist.

20

In another embodiment, the present invention provides 2-naphthamide derivatives of the formula (I'), its tautomeric or stereoisomeric form, or a salt thereof:

25



wherein

-R<sup>1</sup> represents -O-R<sup>10</sup>-OR<sup>11</sup>, -OR<sup>11</sup>, -SR<sup>11</sup>, -SOR<sup>11</sup>, -SO<sub>2</sub>R<sup>11</sup>, -NR<sup>12</sup>R<sup>13</sup>, or -CHR<sup>14</sup>R<sup>15</sup>,

wherein

-R<sup>10</sup>- represents (C<sub>1-6</sub>) alkylene;

R<sup>11</sup> represents aryl, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkyl optionally substituted by (C<sub>3-8</sub>)cycloalkyl, aryl or heterocycle comprising 4-9 carbons and at least one N, O, or S as a heteroatom

wherein

said (C<sub>3-8</sub>)cycloalkyl, aryl and heterocycle optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, (C<sub>1-6</sub>) alkyl optionally substituted by mono-, di-, or tri halogen, and (C<sub>1-6</sub>) alkoxy optionally substituted by (C<sub>3-8</sub>)cycloalkyl, or mono-, di-, or tri halogen;

R<sup>12</sup> and R<sup>13</sup> independently represent hydrogen, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkyl optionally substituted by aryl or heteroaryl,

or

R<sup>12</sup> and R<sup>13</sup> form, together with the nitrogen atom, a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

R<sup>14</sup> and R<sup>15</sup> independently represent hydrogen, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally

- 10 -

substituted by aryl or heteroaryl, (C<sub>1-6</sub>) alkyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkoxy optionally substituted by aryl or heteroaryl,

or

5

R<sup>14</sup> and R<sup>15</sup> form, together with the CH, a (C<sub>3-8</sub>)cycloalkyl optionally interrupted by NH, or O, or a phenyl optionally substituted by hydroxy, halogen or (C<sub>1-6</sub>) alkyl;

10

R<sup>21</sup> represents hydroxy, cyano, amino, (C<sub>1-6</sub>)alkylamino, thienyl, pyridyl, naphthyl, 1H-pyrrolo[2,3-b]pyridin-3-yl, indolyl optionally substituted by halogen or hydroxy, or phenyl,

15

wherein said phenyl and naphthyl optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, amino, N((C<sub>1-6</sub>) alkyl)amino, di(C<sub>1-6</sub>) alkylamino, N((C<sub>1-6</sub>) alkyl sulfonyl)amino, morpholino, phenyl, pyridyl, (C<sub>1-6</sub>) alkoxy optionally substituted by mono-, di-, or tri halogen, and (C<sub>1-6</sub>) alkyl optionally substituted by mono-, di-, or tri halogen; and

20

R<sup>22</sup> represents hydrogen or hydroxy.

Yet another embodiment of the compounds of formula (I) or (I') are those wherein:

25

R<sup>1</sup> represents phenoxy, (C<sub>1-6</sub>) alkoxy optionally substituted by cyclopropyl, cyclohexyl, pyrrolidinyl, piperidinyl, imidazolyl, pyridyl, pyrrolyl, phenyl, or thiazolyl optionally substituted by (C<sub>1-6</sub>)alkyl,

wherein

30

said phenyl has optionally one or two substituents selected from the group consisting of fluoro, chloro, bromo, nitro, hydroxy, (C<sub>1-6</sub>)alkyl optionally substituted by mono-, di, or tri halogen, and (C<sub>1-6</sub>) alkoxy optionally substituted by mono-, di, or tri halogen, cyclopropyl, or cyclohexyl.

Another embodiment of the compounds of formula (I) or (I') are those wherein:

R<sup>1</sup> represents phenoxy(C<sub>1-6</sub>)alkyl, phenoxy(C<sub>1-6</sub>)alkenyl, phenoxy(C<sub>1-6</sub>)alkynyl, or phenyl(C<sub>1-6</sub>)alkoxy.

Further embodiment of the compounds of formula (I) is those

wherein

R<sup>2</sup> represents phenyl (C<sub>1-6</sub>)alkyl,

wherein

said phenyl has optionally one or two substituents selected from the group consisting of fluoro, chloro, bromo, iodo, hydroxy, nitro, amino, N(methanesulfonyl)amino, morpholino, phenyl, pyridyl, methoxy, ethoxy, and trifluoromethyl.

Further embodiment of the compounds of formula (I') is those wherein:

R<sup>1</sup> represents phenoxy, (C<sub>1-6</sub>) alkoxy optionally substituted by cyclopropyl, cyclohexyl, pyrrolidinyl, piperidinyl, imidazolyl, pyridyl, pyrrolyl, phenyl, or thiazolyl optionally substituted by (C<sub>1-6</sub>)alkyl,

wherein

said phenyl has optionally one or two substituents selected from the group consisting of fluoro, chloro, bromo, nitro, hydroxy, (C<sub>1-6</sub>)alkyl optionally substituted by mono-, di, or tri halogen, and (C<sub>1-6</sub>) alkoxy optionally substituted by mono-, di, or tri halogen, cyclopropyl, or cyclohexyl;

R<sup>21</sup> represents cyano, thienyl, pyridyl, phenyl, naphthyl, 1H-pyrrolo[2,3-b]pyridin-3-yl, or indolyl optionally substituted by halogen or hydroxy,

wherein

said phenyl and naphthyl have one or two substituents selected from the group consisting of fluoro, chloro, bromo, hydroxy, nitro, amino, N((C<sub>1-6</sub>) alkyl)amino, di(C<sub>1-6</sub>) alkylamino, N((C<sub>1-6</sub>) alkyl sulfonyl)-amino, morpholino, phenyl, pyridyl, trifluoromethyl, trifluoromethyloxy, (C<sub>1-6</sub>) alkoxy, and (C<sub>1-6</sub>) alkyl; and

R<sup>22</sup> represents hydrogen or hydroxy.

More preferably, said 2-naphthamide derivatives of the formula (I) or (I') are selected from the group consisting of:

N-[6-(benzyloxy)-2-naphthoyl]phenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-(trifluoromethyl)phenylalanine;  
N-{6-[(4-fluorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-fluorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(2-fluorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-[6-(3-pyridinylmethoxy)-2-naphthoyl]phenylalanine;  
N-{6-[(3,4-difluorobenzyl)oxy]-2-naphthoyl}phenylalanine;

- N-{6-[2-(1H-pyrrol-1-yl)ethoxy]-2-naphthoyl}phenylalanine;  
N-[6-(4-pyridinylmethoxy)-2-naphthoyl]phenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(trifluoromethyl)phenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]tryptophan;  
5 N-[6-(benzyloxy)-2-naphthoyl]-O-methyltyrosine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-methoxytyrosine;  
N-[6-(benzyloxy)-2-naphthoyl]- $\beta$ -hydroxyphenylalanine;  
N-[6-(2-phenylethoxy)-2-naphthoyl]phenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-chlorophenylalanine;  
10 N-[6-(benzyloxy)-2-naphthoyl]-3-fluorophenylalanine;  
N-{6-[(2-chlorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-chlorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(2-methoxybenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-methoxybenzyl)oxy]-2-naphthoyl}phenylalanine;  
15 N-{6-[(2,3-dichlorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3,5-dichlorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3,5-dimethoxybenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(2-thienyl)alanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-bromophenylalanine;  
20 N-[6-(benzyloxy)-2-naphthoyl]-4-nitrophenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-hydroxyphenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(1-naphthyl)alanine;  
N-[6-(benzyloxy)-2-naphthoyl]-5-hydroxytryptophan;  
N-[6-(benzyloxy)-2-naphthoyl]-2-fluorophenylalanine;  
25 N-{6-[(2-bromobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-bromobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(2-methylbenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-methylbenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-nitrobenzyl)oxy]-2-naphthoyl}phenylalanine;  
30 N-[6-(benzyloxy)-2-naphthoyl]-3-(2-naphthyl)alanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-iodophenylalanine;

- N-[6-(benzyloxy)-2-naphthoyl]-5-fluorotryptophan;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(1H-pyrrolo[2,3-b]pyridin-3-yl)alanine;  
N-{6-[2-(4-pyridinyl)ethoxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-ethoxybenzyl)oxy]-2-naphthoyl}phenylalanine; and  
5 N-[6-(2-phenylpropoxy)-2-naphthoyl]phenylalanine;

Further, the present invention provides a medicament which include one of the compounds described above and optionally pharmaceutically acceptable excipient.

- 10 The Alkyl per se and "alk" and "alkyl" in alkoxy, alkylene, alkanoyl, alkylamino, alkylaminocarbonyl, alkylaminosulphonyl, alkylsulphonylamino, alkoxycarbonyl, alkoxycarbonylamino and alkanoylamino represent a linear or branched alkyl radical having generally 1 to 6, preferably 1 to 4 and particularly preferably 1 to 3 carbon atoms, representing illustratively and preferably methyl, ethyl, n-propyl, isopropyl,  
15 tert-butyl, n-pentyl and n-hexyl.

Alkoxy illustratively and preferably represents methoxy, ethoxy, n-propoxy, isopropoxy, tert-butoxy, n-pentoxy and n-hexoxy.

- 20 Alkylamino represents an alkylamino radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylamino, ethylamino, n-propylamino, isopropylamino, tert-butylamino, n-pentylamino, n-hexyl-amino, N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, N-methyl-N-n-propylamino, N-isopropyl-N-n-propylamino, N-t-butyl-N-methylamino,  
25 N-ethyl-N-n-pentylamino and N-n-hexyl-N-methylamino.

Aryl per se represents a mono- to tricyclic aromatic carbocyclic radical having generally 6 to 14 carbon atoms, illustratively and preferably representing phenyl, naphthyl and phenanthrenyl.




Heterocyclic ring per se and hetero ring in heteroaryl refers to a 3- to 15-membered ring radical which consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. The heterocyclic ring radical may be a monocyclic, bicyclic or tricyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclic ring radical may be optionally oxidized and the heterocyclic ring system may be partially or fully saturated or aromatic. Examples of such rings include, but are not limited to thienyl, furyl, benzothienyl, furanyl, benzofuranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, pyrimidinyl, pyrrolyl, isothiazolyl, thiazolyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl, imidazolyl, thiadiazolyl, benzothiadiazolyl, oxadiazolyl, benzothiazolyl, indolyl, indazolyl, carbazolyl, quinolyl, isoquinolyl, benzodioxolyl, indazolyl, indazolinolyl, pyrrolidinyl, piperidinyl, pyranyl, pyrazolinyl, piperazinyl, morpholinyl, thiamorpholinyl, thiazolidinyl, benzofuranoyl, thiamorpholinyl sulfone, benzoxazolyl, oxopiperidinyl, oxopyrrolidinyl, oxoazopinyl, azepinyl, furazanyl, tetrahydropyranyl, tetrahydrofuranyl, dioxolyl, dioxinyl, oxathieryl, benzodioxolyl and the like

#### EMBODIMENT OF THE INVENTION

The compound of the formula (I) of the present invention can be, but not limited to be, prepared by combining various known methods. In some embodiments, one or more of the substituents, such as amino group, carboxyl group, and hydroxyl group of the compounds used as starting materials or intermediates are advantageously protected by a protecting group known to those skilled in the art. Examples of the protecting groups are described in "Protective Groups in Organic Synthesis (3rd Edition)" by Greene and Wuts, John Wiley and Sons, New York 1999.

The compound of the formula (I) of the present invention can be, but not limited to be, prepared by the method [A] below.

## Method [A]




- 5 The compound of the formula (I) (wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $m$ , and  $n$  are the same as defined above) can be prepared by deprotection of the compound of formula (II) (wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $m$ , and  $n$  are the same as defined above and  $X$  represents  $C_{1-6}$  alkyl, benzyl, 4-methoxybenzyl or 3,4-dimethoxybenzyl).
- 10 The deprotection of carboxyl group can be conducted by using a base including, for instance, an alkali metal alkoxide such as sodium methoxide, sodium ethoxide and potassium tert-butoxide; alkali metal hydroxide such as sodium hydroxide, lithium hydroxide and potassium hydroxide, or an acid including, for instance, HCl, HBr, trifluoroacetic acid and  $BBr_3$ .
- 15 The removal of protective group  $Z_1$  can be conducted by using a base including, for instance, sodium hydroxide, lithium hydroxide and potassium hydroxide, or an acid including, for instance, HCl, HBr, trifluoroacetic acid and  $BBr_3$ .
- 20 The deprotection can also be done by hydrogenation using a catalyst including, for instance, palladium on carbon and palladium hydroxide, when  $Z_1$  is benzyl, 4-methoxybenzyl or 3,4-dimethoxybenzyl.
- 25 Also, the deprotection can be done by using a reagent such as ceric ammonium nitrate (CAN) or 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ), when  $Z_1$  is 4-methoxybenzyl or 3,4-dimethoxybenzyl.

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); alcohols such as methanol, ethanol, 1-propanol, isopropanol and tert-butanol; water, and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 100°C. The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

#### Preparation of compound formula (II)

##### Procedure A-I



The compound of formula (II) (wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $X$ ,  $m$  and  $n$  are the same as defined above) can be prepared by the reaction of compound (III) (wherein  $R^1$  is the same as defined above) or (III') (wherein  $R^1$  is the same as defined above and  $L_1$  is leaving group, for instance, halogen atom such as chlorine, bromine, or iodine atom

and azole such as 1,3-imidazole and 1,2,4-triazole) with the compound of formula (IV) (wherein  $R^2$ ,  $R^3$ , X, m and n are the same as defined above).

5 The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); and others.  
10 Optionally, two or more of the solvents selected from the listed above can be mixed and used.


The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 0°C to 100°C.  
15 The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

The reaction of compound of (III) and (IV) may be carried out using coupling agent including, for instance, carbodiimides such as N, N-dicyclohexylcarbodiimide and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, 1-hydroxybenzotriazole, and others.  
20

The reaction of compound of (III) and (IV) can also be advantageously conducted in the presence of a base, including, for instance, such as pyridine, triethylamine and N,N-diisopropylethylamine, dimethylaniline, diethylaniline, and others.  
25

The compound (III), (III') and amine (IV) can be commercially available or can be prepared by the use of known techniques.

## Procedure A-II -a



5 Step 1; the compound of formula (IIa) (wherein  $R^2$ ,  $R^3$ ,  $R^{11}$ ,  $X$ ,  $m$  and  $n$  are the same as defined above and  $A$  represents  $O$  or  $S$ ) can be prepared by the reaction of compound (Va) (wherein  $R^2$ ,  $R^3$ ,  $X$ ,  $m$  and  $n$  are the same as defined above and  $A$  represents  $O$  or  $S$ ) with the compound of formula (VIa) (wherein  $R^{11}$  is the same as defined above and  $Y$  represents a leaving group, e.g., halogen, and alkylsulfonyloxy).

10

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as  $N$ ,  $N$ -dimethylformamide (DMF),  $N$ ,  $N$ -dimethylacetamide and  $N$ -methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); ketones such as acetone; alcohols such as methanol, ethanol, 1-propanol, isopropanol and tert-butanol; and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

20

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about  $0^\circ\text{C}$  to  $100^\circ\text{C}$ .

The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

5 The reaction can be advantageously carried out in the presence of a base including, for instance, an alkali metal hydride such as sodium hydride or potassium hydride; alkali metal alkoxide such as sodium methoxide, sodium ethoxide and potassium tert-butoxide; alkali metal hydroxide such as sodium hydroxide and potassium hydroxide; alkali metal carbonates such as sodium carbonate and potassium carbonate; alkali metal hydrogen carbonates such as sodium hydrogen carbonate and  
10 potassium hydrogen carbonate; alkaline earth metal alkoxides such as magnesium ethoxide; organic amines such as pyridine, triethylamine and N,N-diisopropylethylamine, dimethylaniline, diethylaniline and others.

15 Step 1'; the compound of formula (IIa) (wherein  $R^2$ ,  $R^3$ , X, m and n are the same as defined above, A represents O and  $R^{11}$  represents aryl) can be prepared by the reaction of compound (Va) (wherein  $R^2$ ,  $R^3$ , X, m and n are the same as defined above, A represents O and  $R^{11}$  represents aryl) with the compound of formula (VIa') (wherein  $R^{11}$  represents aryl and M represents metal group including, for instance, organoborane group such as boronic acid and di-methoxy boryl; organostannyl group  
20 such as tributyl stannyl, and the like) in the presence of a copper catalyst such as copper(II) acetate and others.

The reaction can be advantageously carried out in the presence of a base including, for instance, cesium carbonate, sodium carbonate, potassium carbonate, pyridine,  
25 triethylamine and others.

The reaction may be carried out in a solvent including, for instance, ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such  
30 as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methyl-

pyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); alcohols such as methanol, ethanol, 1-propanol, isopropanol and tert-butanol; and others.

Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 0°C to 120°C. The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.


Step 2; the compound of formula (VIIIa) (wherein  $R^{11}$  is the same as defined above, A represents O or S and  $Z_1$  represents  $C_{1-6}$  alkyl, benzyl, 4-methoxybenzyl, 3, 4-dimethoxybenzyl and others) can be prepared by the reaction of compound (VIIa) (wherein  $Z_1$  is the same as defined above and A represents O or S) with the compound of formula (VIa) (wherein  $R^{11}$  is the same as defined above and Y represents a leaving group, e.g., halogen, and alkylsulfonyloxy) in the similar manner described in the step 1 of Procedure A-II-a, for the preparation of the compound of formula (IIa).

Step 3; the compound of formula (IIa) (wherein  $R^2$ ,  $R^3$ ,  $R^{11}$ , X, m and n are the same as defined above and A represents O or S) can be prepared by 1) the removal of protective group  $Z_1$  of the compound of the formula (VIIIa) and then 2) the reaction with the compound of the formula (IV) (wherein  $R^2$ ,  $R^3$ , X, m and n are the same as defined above).

The removal of protective group  $Z_1$  can be done in the similar manner described in the Method A for the preparation of the compound of formula (I).

The successive reaction with the compound of the formula (IV) (wherein  $R^2$ ,  $R^3$ , X, m and n are the same as defined above) can be carried out in the similar manner described in Procedure A-I, for the preparation of the compound of formula (II).

# 5 Procedure A-II-b



Step 1; the compound of formula (VIIIb) (wherein  $R^{13}$  is the same as defined above and  $Z_2$  represents  $C_{1-6}$  alkyl, benzyl, 4-methoxybenzyl, 3, 4-dimethoxybenzyl and others) can be prepared by the reaction of compound (VIIb) (wherein  $Z_2$  is the same as defined above) with the compound of formula (VIb) (wherein  $R^{13}$  is the same as defined above and Y represents a leaving group, e.g., halogen, and alkylsulfonyloxy).

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide and N-methylpyrrolidone; sulfoxides such as dimethylsulfoxide (DMSO); ketones such as acetone; alcohols such as methanol, ethanol, 1-propanol, isopropanol and tert-



butanol; and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

5 The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 180°C. The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 2 to 24 hours.

10 The reaction can be advantageously carried out in the presence of a base including, for instance, cesium carbonate, sodium carbonate, potassium carbonate, pyridine, triethylamine and others.

15 Step 2; the compound of formula (IIb-i) (wherein  $R^2$ ,  $R^3$ ,  $R^{13}$ , X, m and n are the same as defined above) can be prepared by 1) the removal of protective group  $Z_2$  of the compound of the formula (VIIIb) and then 2) the reaction with the compound of the formula (IV) (wherein  $R^2$ ,  $R^3$ , X, m and n are the same as defined above) in the similar manner described in Procedure A-I, for the preparation of the compound of formula (II).

20 Step 3, the compound of formula (IIb-ii) (wherein  $R^2$ ,  $R^3$ ,  $R^{12}$ ,  $R^{13}$ , X, m and n are the same as defined above) can be prepared by the reaction of compound (IIb-i) (wherein  $R^2$ ,  $R^3$ ,  $R^{13}$ , X, m and n are the same as defined above) with the compound of formula (VIb') (wherein  $R^{12}$  is the same as defined above and Y represents a leaving group, e.g., halogen and alkylsulfonyloxy) in the similar manner described in  
25 step 1 of Procedure A-II-a, for the preparation of the compound of formula (II-a).

The compound (Va) can be commercially available or can be prepared by either the use of the similar procedure for the preparation of the compound of formula (II) or known techniques. The compound (VIa), (VIa'), (VIIa), (VIb), (VIb') and (VIIb) can  
30 be commercially available or can be prepared by the use of known techniques.

When the compound shown by the formula (I) or a salt thereof has an asymmetric carbon in the structure, their optically active compounds and racemic mixtures are also included in the scope of the present invention.

5 Typical salts of the compound shown by the formula (I) include salts prepared by reaction of the compounds of the present invention with a mineral or organic acid, or an organic or inorganic base. Such salts are known as acid addition and base addition salts, successively.

10 Acids to form salts include inorganic acids such as, without limitation, sulfuric acid, phosphoric acid, hydrochloric acid, hydrobromic acid, hydriodic acid and the like, and organic acids, such as, without limitation, p-toluenesulfonic acid, methane-sulfonic acid, oxalic acid, p-bromophenylsulfonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like.

15 Base addition salts include those derived from inorganic bases, such as, without limitation, ammonium hydroxide, alkaline metal hydroxide, alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, and organic bases, such as, without limitation, ethanolamine, triethylamine, tris(hydroxymethyl)aminomethane,  
20 and the like. Examples of inorganic bases include, sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like.

25 The compound of the present invention or a salts thereof, depending on its substituents, may be modified to form lower alkylesters or known other esters; and/or hydrates or other solvates. Those esters, hydrates, and solvates are included in the scope of the present invention.

30 The compound of the present invention may be administered in oral forms, such as, without limitation normal and enteric coated tablets, capsules, pills, powders, granules, elixirs, tinctures, solution, suspensions, syrups, solid and liquid aerosols

and emulsions. They may also be administered in parenteral forms, such as, without limitation, intravenous, intraperitoneal, subcutaneous, intramuscular, and the like forms, well-known to those of ordinary skill in the pharmaceutical arts. The compounds of the present invention can be administered in intranasal form via  
5 topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal delivery systems well-known to those of ordinary skilled in the art.

The dosage regimen with the use of the compounds of the present invention is selected by one of ordinary skill in the arts, in view of a variety of factors, including,  
10 without limitation, age, weight, sex, and medical condition of the recipient, the severity of the condition to be treated, the route of administration, the level of metabolic and excretory function of the recipient, the dosage form employed, the particular compound and salt thereof employed.

15 The compounds of the present invention are preferably formulated prior to administration together with one or more pharmaceutically-acceptable excipients. Excipients are inert substances such as, without limitation carriers, diluents, flavoring agents, sweeteners, lubricants, solubilizers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

20 Yet another embodiment of the present invention is pharmaceutical formulation comprising a compound of the invention and one or more pharmaceutically-acceptable excipients that are compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Pharmaceutical formulations  
25 of the invention are prepared by combining a therapeutically effective amount of the compounds of the invention together with one or more pharmaceutically-acceptable excipients. In making the compositions of the present invention, the active ingredient may be mixed with a diluent, or enclosed within a carrier, which may be in the form of a capsule, sachet, paper, or other container. The carrier may serve as a diluent,  
30 which may be solid, semi-solid, or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions,

solutions, syrups, aerosols, ointments, containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

5 For oral administration, the active ingredient may be combined with an oral, and non-toxic, pharmaceutically-acceptable carrier, such as, without limitation, lactose, starch, sucrose, glucose, sodium carbonate, mannitol, sorbitol, calcium carbonate, calcium phosphate, calcium sulfate, methyl cellulose, and the like; together with, optionally, disintegrating agents, such as, without limitation, maize, starch, methyl  
10 cellulose, agar bentonite, xanthan gum, alginic acid, and the like; and optionally, binding agents, for example, without limitation, gelatin, natural sugars, beta-lactose, corn sweeteners, natural and synthetic gums, acacia, tragacanth, sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like; and, optionally, lubricating agents, for example, without limitation, magnesium stearate, sodium  
15 stearate, stearic acid, sodium oleate, sodium benzoate, sodium acetate, sodium chloride, talc, and the like.

In powder forms, the carrier may be a finely divided solid which is in admixture with the finely divided active ingredient. The active ingredient may be mixed with a  
20 carrier having binding properties in suitable proportions and compacted in the shape and size desired to produce tablets. The powders and tablets preferably contain from about 1 to about 99 weight percent of the active ingredient which is the novel composition of the present invention. Suitable solid carriers are magnesium carboxymethyl cellulose, low melting waxes, and cocoa butter.

25 Sterile liquid formulations include suspensions, emulsions, syrups and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent, or a mixture of both sterile water and sterile organic solvent.

The active ingredient can also be dissolved in a suitable organic solvent, for example, aqueous propylene glycol. Other compositions can be made by dispersing the finely divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

5

The formulation may be in unit dosage form, which is a physically discrete unit containing a unit dose, suitable for administration in human or other mammals. A unit dosage form can be a capsule or tablets, or a number of capsules or tablets. A "unit dose" is a predetermined quantity of the active compound of the present invention, calculated to produce the desired therapeutic effect, in association with one or more excipients. The quantity of active ingredient in a unit dose may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved.

10

Typical oral dosages of the present invention, when used for the indicated effects, will range from about 0.01 mg/kg/day to about 100 mg/kg/day, preferably from 0.1 mg/kg/day to 30 mg/kg/day, and most preferably from about 0.5 mg/kg/day to about 10 mg/kg/day. In the case of parenteral administration, it has generally proven advantageous to administer quantities of about 0.001 to 100 mg/kg/day, preferably from 0.01 mg/kg/day to 1 mg/kg/day. The compounds of the present invention may be administered in a single daily dose, or the total daily dose may be administered in divided doses, two, three, or more times per day. Where delivery is via transdermal forms, of course, administration is continuous.

15  
20

### Examples

5 The present invention will be described in detail below in the form of examples, but they should by no means be construed as defining the metes and bounds of the present invention.

In the examples below, all quantitative data, if not stated otherwise, relate to percentages by weight.

10 Melting points are uncorrected. Liquid Chromatography - Mass spectroscopy (LC-MS) data were recorded on a Micromass Platform LC with Shimadzu Phenomenex ODS column (4.6 mm x 30 mm) flushing a mixture of acetonitrile-water (9:1 to 1:9) at 1 ml/min of the flow rate. Mass spectra were obtained using  
15 electrospray (ES) ionization techniques (micromass Platform LC). TLC was performed on a precoated silica gel plate (Merck silica gel 60 F-254). Silica gel (WAKO-gel C-200 (75-150  $\mu$ m)) was used for all column chromatography separations. All chemicals were reagent grade and were purchased from Sigma-Aldrich, Wako pure chemical industries, Ltd., Great Britain, Tokyo kasei kogyo Co., Ltd., Japan, Nacalai tesque, Inc., Watanabe Chemical Ind. Ltd., Maybridge plc,  
20 Lancaster Synthesis Ltd., Great Britain, Merck KGaA, Germany, Kanto Chemical Co., Ltd.

The effect of the present compounds were examined by the following assays and pharmacological tests.

25

[Measurement of the [ $^3$ H]-iloprost binding to HEL cells] (Assay 1)

A human erythroleukemia cell line, HEL 92.1.7, was purchased from American Type Culture Collection and maintained in RPMI-1640 medium (Gibco BRL)  
30 supplemented with 10% fetal calf serum (FCS), 2 mM glutamine, 4.5 g/L glucose, 10 mM Hepes, 1 mM sodium pyruvate, 100 U/ml penicillin, and 100  $\mu$ g/ml strepto-

mycin in a humidified 5% CO<sub>2</sub> atmosphere at 37°C. Cells were collected with centrifugation and washed with binding assay buffer (BAB: 50 mM Tris-HCl, 5 mM MgCl<sub>2</sub> (pH 7.5)). Cells were suspended at the density of 6.25 x 10<sup>6</sup> cells/ml in BAB, and one million cells in 160 µl aliquot of cell suspension were put in a well of 96 well plate (Falcon). Then, 20 µl of compound solution, 100 µM of iloprost (for non-specific binding), or buffer alone (total binding), diluted with 1% DMSO in BAB was added. Finally, another 20 µl containing [<sup>3</sup>H]-iloprost (0.02 µCi, 0.5-1 pmol) in BAB was added and incubated at room temperature for 30 min with a gentle shaking. Cell suspension was then transferred to a well of MultiScreen plate with GF/C glass filters (Millipore) to harvest cells. Cells were washed twice with 200 µl of ice-cold BAB and the plate was kept at 55°C for 30 min to dry filters. The filter in the well was punched out to a counting tube and 2 ml of Ultima Gold XR (Packard) was added. [<sup>3</sup>H]-radio activity in the filter was measured by a liquid scintillation counter (Beckman, USA).

#### [Iloprost-induced cAMP production assay in HEL cells] (Assay 2)

HEL cells were collected with centrifugation and washed with cAMP assay buffer (CAB: Hank's balanced salt solution, 17 mM Hepes, 0.1% bovine serum albumin, 1 mM IBMX, 0.4% DMSO, and 1 mM L-ascorbic acid sodium salt (pH 7.4)). Cells were suspended at the density of 2.5 x 10<sup>5</sup> cells/ml in CAB, and twenty thousand cells in 80 µl aliquot of cell suspension were put in a well of 96 well plate (Falcon). Then, 10 µl of compound solution diluted with 1% DMSO in CAB or buffer alone was added. The plate was incubated at 37°C for 30 min. Then, another 10 µl containing 100 nM iloprost in CAB or buffer alone was added and further incubated at 37°C for 30 min. cAMP content in the well was measured by a cAMP ELISA kit (Applied Biosystems, USA).

[Measurement of rhythmic bladder contraction in anesthetized rats]

(1) Animals

5 Female Sprague-Dawley rats (200~250 g / Charles River Japan) were used.

(2) Rhythmic bladder contraction in anesthetized rats

10 Rats were anesthetized by intraperitoneal administration of urethane (Sigma) at 1.25 g/kg. The trachea was cannulated with a polyethylene tube (HIBIKI, No. 8) to facilitate respiration; and a cannula (BECTON DICKINSON, PE-50) was placed in the left femoral vein for intravenous administration of testing compounds. The abdomen was opened through a midline incision, and after both ureters were cut, a water-filled balloon (about 1 ml capacity) was inserted through the apex of the bladder dome. The balloon was connected to a pressure transducer onto a polygraph. Rhythmic bladder contraction was elicited by raising up intravesical pressure to approximately 15 cm H<sub>2</sub>O. After the rhythmic bladder contraction was stable, a testing compound was administered intravenously. Activity was estimated by measuring disappearance time and amplitude of the rhythmic bladder contraction. 20 The effect on amplitude of bladder contractions was expressed as a percent suppression of the amplitude of those after the disappearance was recovered. Experimental values were expressed as the mean±S.E.M. The testing compounds-mediated inhibition of the rhythmic bladder contraction was evaluated using Student's t-test. A probability level less than 5% was accepted as significant difference. 25

Results of Iloprost-induced cAMP production assay (Assay2) are shown in Examples and tables of the Examples below. For practical reasons, the compounds are grouped in three classes of activity as follows:

30


A (= or <) 0.1  $\mu$ M < B (= or <) 1  $\mu$ M < C



The compounds of the present invention also show excellent selectivity, and strong activity in vivo assays.

5 **Example 1-1**

(1) N-(6-Hydroxy-2-naphthoyl)phenylalanine Methyl Ester




10

15

To a solution of 6-hydroxy-2-naphthoic acid (300 mg), DL-phenylalanine methyl ester (344 mg), 1-hydroxybenzotriazole (HOBt, 280mg), and triethylamine (0.3 ml) in N,N-dimethylformamide (DMF, 8 ml) was added 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide (EDCI, 396 mg). The mixture was stirred at room temperature overnight and concentrated in vacuo. The residue was extracted with ethyl acetate and washed with brine. The organic layer was dried over sodium sulfate and evaporated to give colorless viscous oil, that was purified by silica gel column chromatography (hexane/ethyl acetate = 1:1) to give N-(6-hydroxy-2-naphthoyl)phenylalanine methyl ester (475 mg, 85%) as a colorless foam.


20

(2) N-(6-Benzyloxy-2-naphthoyl)phenylalanine Methyl Ester



To a solution of N-(6-hydroxy-2-naphthoyl)phenylalanine methyl ester (200 mg) and benzyl chloride (80  $\mu$ l) in DMF (5 ml) was added potassium carbonate (95 mg). The mixture was stirred at room temperature overnight and at 70°C for 2 hours. The solvent was evaporated in vacuo and the residue was extracted with ethyl acetate and washed with brine. The organic layer was dried over sodium sulfate and concentrated in vacuo to give a colorless solid, that was purified by preparative TLC (chloroform/ethyl acetate = 10/1) and crystallization with diisopropyl ether to give N-(6-benzyloxy-2-naphthoyl)phenylalanine methyl ester (190 mg, 76%) as a colorless solid.

(3) N-(6-Benzyloxy-2-naphthoyl)phenylalanine



To a solution of N-(6-benzyloxy-2-naphthoyl)phenylalanine methyl ester (60 mg) in ethanol (2 ml) was added 1N lithium hydroxide water solution (0.2 ml). The mixture was stirred at 50°C for 4 hours and concentrated in vacuo. The residue was diluted with water and acidified with 1N hydrochloric acid to be extracted with a mixture of acetate and tetrahydrofurane. The organic layer was washed with brine and dried over sodium sulfate. The solvent was evaporated off to give a colorless oil, that was crystallized with diisopropylether to give N-(6-benzyloxy-2-naphthoyl)phenylalanine (47.8 mg, 82%) as a colorless solid.

Melting point: 230°C

Molecular weight: 425.48

Mass spectrometry: 426







Activity grade assay 2: A







- 33 -






<sup>1</sup>H-NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 3.11 (1H, dd, J = 13.9, 10.4 Hz), 3.22 (1H, dd, J = 13.8, 4.4 Hz), 4.64-4.69 (1H, m), 5.25 (2H, s), 7.17 (1H, t, J = 7.2 Hz), 7.25-7.35 (6H, m), 7.42 (2H, t, J = 7.2 Hz), 7.48 (1H, d, J = 2.3 Hz), 7.52 (2H, d, J = 7.3 Hz),  
5 7.84 (2H, d, J = 0.9 Hz), 7.93 (1H, d, J = 9.1 Hz), 8.34 (1H, s), 8.73 (1H, d, J = 8.2 Hz), 12.76 (1H, bs).







In the similar manner as described in Example 1-1, compounds in Example 1-2 to 1-94 as shown in Table 1 were synthesized.







**Table 1**

Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-2		365,39	366	196-198	C
1-3		493,49	494	212-214 (dec.)	A
1-4		455,52	456	225-227	B
1-5		443,48	444	197	A
1-6		443,48	444	197-199	A
1-7		443,48	444	180-182	A







Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-8		426,48	427	217	A
1-9		426,48	427	214	B
1-10		461,47	462	185	A
1-11		428,49	429	168-169	A
1-12		426,48	427	220-221	A
1-13		493,49	494	ND	B

Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-14		493,49	494	ND	A
1-15		464,53	465	ND	A
1-16		455,52	456	ND	A
1-17		471,51	472	ND	A
1-18		441,49	442	ND	A







Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-19		439,52	440	144	A
1-20		459,93	460	202-204	A
1-21		443,48	444	188-191	A
1-22		443,48	444	214-216	A
1-23		459,93	460	215	A
1-24		459,93	460	187	A







Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-25		459,93	460	226	B
1-26		493,49	494	167	B
1-27		493,49	494	158	C
1-28		493,49	494	224	C
1-29		455,52	456	169	A
1-30		455,52	456	163	A















Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-31		455,52	456	188	C
1-32		494,38	494 (M)	202	B
1-33		494,38	494 (M)	211	A
1-34		494,38	494 (M)	180	A
1-35		485,54	486	140	A
1-36		509,49	510	157	B







Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-37		509,49	510	158	C
1-38		485,54	486	161	C
1-39		484,51	485	202	B
1-40		453,54	454	210	B
1-41		494,38	494 (M)	245	B
1-42		494,38	494 (M)	202	C







Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-43		426,48	427	172-173 (dec.)	B
1-44		431,51	432	200-202	A
1-45		504,38	505	219-220	A
1-46		470,49	469 (M-1)	227-228	A
1-47		441,49	442	214	A
1-48		475,55	476	218	A

Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-49		480,53	481	261-262 (dec.)	A
1-50		443,48	444	211-213	A
1-51		504,38	504 (M)	206-207	A
1-52		504,38	504	194	A
1-53		504,38	504	232	C
1-54		439,52	440	204	A







Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-55		439,52	440	184-185	A
1-56		439,52	440	212	B
1-57		453,54	454	167-168	B
1-58		453,54	454	189	B
1-59		461,47	462	199	B
1-60		439,52	440	170-172	B

Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-61		470,49	488	193-194	A
1-62		446,53	447	208	B
1-63		442,52	443	195	B
1-64		453,54	454	182	B
1-65		391,47	392	209	B
1-66		440,50	441	238-240	C





Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-67		475,55	476	226-227	A
1-68		518,59	519	236-237	C
1-69		426,48	427	260-262 (dec.)	B
1-70		475,55	476	220	B
1-71		551,39	552	219-221	A
1-72		439,52	440	223-225	C


Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-73		501,59	502	250-252	B
1-74		431,54	432	220	B
1-75		502,58	503	247-249 (dec.)	B
1-76		510,60	511	216-218 (dec.)	C
1-77		440,50	441	113	B
1-78		440,50	441	143	B



Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-79		482,52	483	236-238 (dec.)	A
1-80		465,51	466	275-277 (dec.)	A
1-81		432,52	433	234	C
1-82		446,55	447	215	C
1-83		440,50	441	190	A
1-84		439,52	440	184	B

Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-85		441,49	442	157	A
1-86		429,48	430	172 (dec.)	B
1-87		442,48	443	252 (dec.)	C
1-88		537,66	538	180	C
1-89		495,58	496	179	B
1-90		497,60	498	172	B


Ex No	Structure	Mol weight	MASS (M+1)	MP	Assay 2
1-91		469,54	470	145	A
1-92		483,57	484	173	B
1-93		497,60	498	140	B
1-94		453,54	454	160	A

**Example 2-1****(1) Benzyl 6-(benzyloxy)-2-naphthoate**

To a solution of 6-hydroxy-2-naphthoic acid (0.50 g, 2.66 mmol) and benzylchloride (1.01 g, 7.97 mmol) in *N,N*-dimethylformamide (15 mL) was added potassium carbonate (1.10 g, 7.97 mmol) and sodium iodide (0.12 g, 0.80 mmol), and the mixture was stirred at 70°C for 4 hours. The mixture was concentrated in vacuo, and the residue was partitioned between ethyl acetate and water. The separated organic phase was dried over sodium sulfate, filtered and concentrated in vacuo. The residue was purified by column chromatography on silica-gel (n-hexane:ethyl acetate, 4:1) to give benzyl 6-(benzyloxy)-2-naphthoate (1.01 g, 103 %) as yellowish granules.

10

15

**(2) 6-(Benzyloxy)-2-naphthoic acid**


To a solution of benzyl 6-(benzyloxy)-2-naphthoate (1.01 g, 2.75 mmol) in ethanol (20 ml) was added dropwise 1N NaOH (5.50 ml, 5.50 mmol) and the mixture was stirred at room temperature for 3 days. The mixture was concentrated under reduced pressure. The residue was diluted with water (10 ml) and washed with ether. The aqueous layer was separated and neutralized with 1N HCl (5.50 ml). The resultant precipitate was collected by filtration, washed successively with water and ethyl

25

acetate and dried under reduced pressure to give 6-(benzyloxy)-2-naphthoic acid (0.42 g, 55 %) as colorless powder.

(3) *tert*-Butyl N-[6-(benzyloxy)-2-naphthoyl]-D-phenylalaninate

5




To a solution of 6-(benzyloxy)-2-naphthoic acid (200 mg, 0.72 mmol), D-phenylalanine *tert*-butyl ester hydrochloride salt (185 mg, 0.72 mmol), 1-hydroxybenzotriazole (280 mg, 0.93 mmol) and triethylamine (0.30 ml, 0.86 mmol) in *N,N*-dimethylformamide (8.0 ml) was added 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (396 mg, 0.93 mmol). The mixture was stirred at room temperature overnight and concentrated in vacuo. The residue was extracted with ethyl acetate and washed with brine. The organic layer was dried over sodium sulfate and evaporated to give colorless viscous oil, that was purified by silica gel column chromatography (hexane:ethyl acetate, 5:1) to give *tert*-butyl N-[6-(benzyloxy)-2-naphthoyl]-D-phenylalaninate (318 mg, 92%) as a colorless solid.

10

15

(4) *N*-[6-(benzyloxy)-2-naphthoyl]-D-phenylalanine

20



*tert*-Butyl *N*-[6-(benzyloxy)-2-naphthoyl]-D-phenylalaninate (100 mg, 0.21 mmol) was dissolved in 4N hydrochloride-dioxane (2.0 ml) and the resulting mixture was kept at room temperature overnight. The mixture was concentrated in vacuo. The residue was triturated with diisopropyl ether to give powder, which was recrystallized from a mixture of methylenechloride and diisopropyl ether to give *N*-[6-(benzyloxy)-2-naphthoyl]-D-phenylalanine (62 mg, 70%) as colorless powder.

Melting point: 179°C

Molecular weight: 425.48

Mass spectrometry: 426

Activity grade assay 2: A

HPLC analysis

Column: Daicel Chiralcel OD-RH 5 $\mu$ m 0.46 cm x 15 cm

Eluent: 0.1% acetic acid in water : acetonitrile = 50 : 50


Flow rate: 1.0 ml / min.

Absorbance: 210 nm

Retention time: 12.29 min. (*N*-[6-(benzyloxy)-2-naphthoyl]-D-phenylalanine)

### Example 2-2

*N*-[6-(benzyloxy)-2-naphthoyl]-L-phenylalanine



According to the similar synthetic procedure as in example 2-1, *N*-[6-(benzyloxy)-2-naphthoyl]-L-phenylalanine was obtained as a colorless powder.

Melting point: 207°C

Molecular weight: 425.48

Mass spectrometry: 426

Activity grade assay2: C

5

#### HPLC analysis

Column: Daicel Chiralcel OD-RH 5µm 0.46 cm x 15 cm

Eluent: 0.1% acetic acid in water : acetonitrile = 50 : 50

Flow rate: 1.0 ml / min.

10


Absorbance: 210 nm

Retention time: 11.10 min. (*N*-[6-(benzyloxy)-2-naphthoyl]-L-phenylalanine)

#### Example 3-1

15

#### *N*-(6-phenoxy-2-naphthoyl)phenylalanine



20

To a mixture of phenylboronic acid (0.141 g, 1.14 mmol), methyl *N*-(6-hydroxy-2-naphthoyl)phenylalaninate (0.200 g, 0.57 mmol), copper(II) acetate (0.104 g, 0.57 mmol) and molecular sieves 4A (570 mg) in dichloromethane was added triethylamine (0.290 g, 2.86 mmol), and stirring was continued at room temperature overnight. The reaction mixture was filtered and concentrated under reduced pressure. The crude product was purified by preparative TLC (hexane: ethyl acetate, 3:1) to give methyl *N*-(6-phenoxy-2-naphthoyl)phenylalaninate as a colorless oil.

25

The methyl ester above was treated with lithium hydroxide monohydrate (0.036 g, 0.86 mmol) in tetrahydrofuran (1.0 mL), ethanol (0.5 mL) and water (1.0 mL) at

60°C for 2 hours. The mixture was neutralized with 1N hydrochloric acid and extracted with ethyl acetate. The separated organic phase was washed with brine, dried over sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by recrystallization (ethyl acetate-hexane) to give *N*-(6-phenoxy-2-naphthoyl)phenylalanine (0.020 g, 9%) as a white powder.

Melting point: 179-180 °C

Molecular weight: 411.46


Mass spectrometry: 412

Activity grade assay 2: C

<sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>): δ 3.11(1H, dd, J = 10.4, 13.9 Hz), 3.22 (1H, dd, J = 4.7, 13.9 Hz), 4.66 (1H, br), 7.10-7.29 (6H, m), 7.30-7.38 (4H, m), 7.46 (2H, m), 7.85 (2H, t, J = 8.8 Hz), 8.06 (1H, d, J = 8.8 Hz), 8.40 (1H, s), 8.77 (1H, br), 12.78 (1H, br).

#### Example 4-1

##### (1) Benzyl 6-(benzylamino)-2-napthoate




To a solution (40 mL) of 6-amino-2-napthoic acid (0.500g, 2.67 mmol) in DMF were added benzyl bromide (1.142 g, 6.68 mmol), potassium carbonate (1.107 g, 8.01 mmol) and sodium iodide (1.201 g, 8.01 mmol), and the mixture was stirred at 80°C overnight. After cooled to room temperature, the mixture was partitioned between ethyl acetate and saturated aqueous ammonium chloride solution. The separated organic phase was washed with water and brine, dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by




column chromatography on silica gel (hexane: ethyl acetate, 3:1) to give benzyl 6-(benzylamino)-2-naphthoate (0.214 g, 21%) as a white solid.

(2) 6-(benzylamino)-2-naphthoic acid



To a solution of benzyl 6-(benzylamino)-2-naphthoate (0.200 g, 0.54 mmol) in tetrahydrofuran (2 mL), ethanol (1 mL) and water (2 mL) was added lithium hydroxide monohydrate (0.068 g, 1.63 mmol), and the mixture was stirred at 60°C overnight. After cooled to room temperature, the mixture was partitioned between ethyl acetate and saturated ammonium chloride aqueous solution. The separated organic phase was washed with water and brine, dried over sodium sulfate, filtered and concentrated under reduced pressure to give 6-benzylamino-2-naphthoic acid (0.132 g, 88%) as a white solid.

(3) Methyl N-[6-(benzylamino)-2-naphthoyl]phenylalaninate




A mixture of 6-(benzylamino)-2-naphthoic acid (0.036 g, 0.13 mmol), DL-phenylalanine hydrochloride salt (0.028 g, 0.13 mmol), EDCI (0.032 g, 0.17 mmol), 1-hydrobenzotriazole (0.023 g, 0.17 mmol) and triethylamine (0.017 g, 0.16 mmol) in DMF (0.6 mL) was stirred at room temperature overnight. The mixture was partitioned between water and ethyl acetate. The separated organic phase was washed

with brine, dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by preparative TLC (dichloromethane: methanol, 10:1) to give methyl N-[6-(benzylamino)-2-naphthoyl]phenylalaninate (0.038 g, 69%) as a white powder.

5

(4) N-[6-(benzylamino)-2-naphthoyl]phenylalanine



10 To a solution of methyl N-[6-(benzylamino)-2-naphthoyl]phenylalaninate (0.038 g, 0.09 mmol) in tetrahydrofuran (1 mL), ethanol (0.5 mL) and water (1 mL) was added lithium hydroxide monohydrate (0.011 g, 0.26 mmol), and the mixture was stirred at 60°C for 2 hours. After cooled to room temperature, the mixture was neutralized with 1N hydrochloric acid solution (0.26 mL) and the solution was extracted with ethyl acetate. The separated organic phase was washed with water and brine, dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by recrystallization (ethyl acetate-hexane) to give N-[6-(benzylamino)-2-naphthoyl]phenylalanine (0.020 g, 53%) as a white solid.

15

20

Melting point: 172°C

Molecular weight: 424.50

Mass spectrometry: 425

Activity grade assay 2: B


25

<sup>1</sup>H-NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 3.09 (1H, dd, J = 10.1, 13.9 Hz), 3.19 (1H, dd, 4.4, 13.9 Hz), 4.39 (2H, d, 5.7 Hz), 6.22 (1H, m), 6.71 (1H, d, 1.6 Hz), 6.82 (1H, t, 5.7 Hz), 7.10 (1H, dd, J = 2.2, 8.8 Hz), 7.17 (1H, m), 7.21-7.27 (3H, m), 7.28-7.36

(4H, m), 7.41 (2H, d,  $J = 7.3$  Hz), 7.49 (1H, d,  $J = 8.8$  Hz), 7.67 (2H, d,  $J = 9.1$  Hz),  
8.15 (1H, s), 8.53 (1H, d,  $J = 8.2$  Hz), 12.7 (1H, Br).

Claims

- (1) A 2-naphthamide derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof:



wherein

m and n independently represent an integer from 0 to 2;

10  
-R¹ represents -O-R¹⁰-OR¹¹, -OR¹¹, -SR¹¹, -S(O)R¹¹, -S(O)₂R¹¹, -NR¹²R¹³, or -CHR¹⁴R¹⁵,

wherein

15  
-R¹⁰- represents (C₁-₆) alkylene;

20  
R¹¹ represents aryl, (C₂-₆)alkenyl optionally substituted by aryl or heteroaryl, (C₂-₆)alkynyl optionally substituted by aryl or heteroaryl, or (C₁-₆) alkyl optionally substituted by (C₃-₈)-cycloalkyl, aryl or heterocycle comprising 4-9 carbons and at least one N, O, or S as a heteroatom,

wherein

25  
said (C₃-₈)cycloalkyl, aryl and heterocycle optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, (C₁-₆) alkyl optionally substituted by

mono-, di-, or tri halogen, and (C<sub>1-6</sub>) alkoxy optionally substituted by (C<sub>3-8</sub>)cycloalkyl, or mono-, di-, or tri halogen;

5 R<sup>12</sup> and R<sup>13</sup> independently represent hydrogen, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkyl optionally substituted by aryl or heteroaryl,

10 or

R<sup>12</sup> and R<sup>13</sup> form, together with the nitrogen atom, a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

15 R<sup>14</sup> and R<sup>15</sup> independently represent hydrogen, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, (C<sub>1-6</sub>) alkyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkoxy optionally substituted by aryl or heteroaryl,

20 or

R<sup>14</sup> and R<sup>15</sup> form, together with the CH, a (C<sub>3-8</sub>)cycloalkyl optionally interrupted by NH, or O, or a phenyl optionally substituted by hydroxy, halogen or (C<sub>1-6</sub>) alkyl;

25 R<sup>2</sup> represents hydrogen, hydroxy, cyano, (C<sub>1-6</sub>) alkoxy, (C<sub>2-6</sub>)alkenyl, (C<sub>2-6</sub>)alkynyl, (C<sub>3-7</sub>)cycloalkyl, or (C<sub>1-6</sub>) alkyl optionally having one or two substituents selected from the group consisting of hydroxy, amino, (C<sub>1-6</sub>)alkylamino, aryl, and heteroaryl comprising 4-10 carbons and at least one N, O, or S as a heteroatom,

30


- 60 -

wherein

said aryl and heteroaryl optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, amino, N((C<sub>1-6</sub>) alkyl sulfonyl)amino, morpholino, phenyl, pyridyl, (C<sub>1-6</sub>) alkoxy optionally substituted by mono-, di-, or tri halogen, and (C<sub>1-6</sub>) alkyl optionally substituted by mono-, di-, or tri halogen; and

R<sup>3</sup> represents hydrogen, or (C<sub>1-6</sub>) alkyl.

- (2) A 2-naphthamide derivative of the formula (I'), its tautomeric or stereoisomeric form, or a salt thereof:



wherein

-R<sup>1</sup> represents -O-R<sup>10</sup>-OR<sup>11</sup>, -OR<sup>11</sup>, -SR<sup>11</sup>, -S(O)R<sup>11</sup>, -S(O)<sub>2</sub>R<sup>11</sup>, -NR<sup>12</sup>R<sup>13</sup>, or -CHR<sup>14</sup>R<sup>15</sup>,

wherein

-R<sup>10</sup>- represents (C<sub>1-6</sub>) alkylene;

R<sup>11</sup> represents aryl, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkyl optionally substituted by (C<sub>3-8</sub>)-

cycloalkyl, aryl or heterocycle comprising 4-9 carbons and at least one N, O, or S as a heteroatom

wherein

said (C<sub>3-8</sub>)cycloalkyl, aryl and heterocycle optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, (C<sub>1-6</sub>) alkyl optionally substituted by mono-, di-, or tri halogen, and (C<sub>1-6</sub>) alkoxy optionally substituted by (C<sub>3-8</sub>)cycloalkyl, or mono-, di-, or tri halogen;

R<sup>12</sup> and R<sup>13</sup> independently represent hydrogen, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkyl optionally substituted by aryl or heteroaryl,

or

R<sup>12</sup> and R<sup>13</sup> form, together with the nitrogen atom, a 5-7 membered saturated hetero ring optionally interrupted by O or NH;

R<sup>14</sup> and R<sup>15</sup> independently represent hydrogen, (C<sub>2-6</sub>)alkenyl optionally substituted by aryl or heteroaryl, (C<sub>2-6</sub>)alkynyl optionally substituted by aryl or heteroaryl, (C<sub>1-6</sub>) alkyl optionally substituted by aryl or heteroaryl, or (C<sub>1-6</sub>) alkoxy optionally substituted by aryl or heteroaryl,

or

$R^{14}$  and  $R^{15}$  form, together with the CH, a (C<sub>3-8</sub>)cycloalkyl optionally interrupted by NH, or O, or a phenyl optionally substituted by hydroxy, halogen or (C<sub>1-6</sub>) alkyl;

5  $R^{21}$  represents hydroxy, cyano, amino, (C<sub>1-6</sub>)alkylamino, thienyl, pyridyl, phenyl, naphthyl, 1H-pyrrolo[2,3-b]pyridin-3-yl, or indolyl optionally substituted by halogen or hydroxy,

wherein

10

said phenyl and naphthyl optionally have one or two substituents selected from the group consisting of halogen, hydroxy, nitro, amino, N((C<sub>1-6</sub>) alkyl)amino, di(C<sub>1-6</sub>) alkylamino, N((C<sub>1-6</sub>) alkyl sulfonyl)-amino, morpholino, phenyl, pyridyl, (C<sub>1-6</sub>) alkoxy optionally substituted by mono-, di-, or tri halogen, and (C<sub>1-6</sub>) alkyl optionally substituted by mono-, di-, or tri halogen; and

15

$R^{22}$  represents hydrogen or hydroxy.

20 (3) The 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

wherein

25  $R^1$  represents phenoxy, (C<sub>1-6</sub>) alkoxy optionally substituted by cyclopropyl, cyclohexyl, pyrrolidinyl, piperidinyl, imidazolyl, pyridyl, pyrrolyl, thiazolyl optionally substituted by (C<sub>1-6</sub>)alkyl, or phenyl,

wherein

30



said phenyl optionally has one or two substituents selected from the group consisting of fluoro, chloro, bromo, nitro, hydroxy, (C<sub>1-6</sub>)alkyl optionally substituted by mono-, di, or tri halogen, and (C<sub>1-6</sub>)alkoxy optionally substituted by mono-, di, or tri halogen, cyclopropyl, or cyclohexyl.

- (4) The 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

wherein

R<sup>1</sup> represents phenoxy(C<sub>1-6</sub>)alkyl, phenoxy(C<sub>1-6</sub>)alkenyl, phenoxy(C<sub>1-6</sub>)alkynyl, or phenyl(C<sub>1-6</sub>)alkoxy.

- (5) The 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

wherein

R<sup>2</sup> represents phenyl (C<sub>1-6</sub>)alkyl,

wherein

said phenyl optionally has one or two substituents selected from the group consisting of fluoro, chloro, bromo, iodo, hydroxy, nitro, amino, N(methanesulfonyl)amino, morpholino, phenyl, pyridyl, methoxy, ethoxy, and trifluoromethyl.

- (6) The 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 2,

wherein

5           R<sup>1</sup>     represents phenoxy, (C<sub>1-6</sub>) alkoxy optionally substituted by cyclopropyl, cyclohexyl, pyrrolidinyl, piperidinyl, imidazolyl, pyridyl, pyrrolyl, phenyl, or thiazolyl optionally substituted by (C<sub>1-6</sub>)alkyl,

          wherein

10           said phenyl has optionally one or two substituents selected from the group consisting of fluoro, chloro, bromo, nitro, hydroxy, (C<sub>1-6</sub>)alkyl optionally substituted by mono-, di, or tri halogen, and (C<sub>1-6</sub>) alkoxy optionally substituted by mono-, di, or tri halogen, cyclopropyl, or cyclohexyl;

15           R<sup>21</sup>   represents cyano, thienyl, pyridyl, phenyl, naphthyl, 1H-pyrrolo[2,3-b]pyridin-3-yl, or indolyl optionally substituted by halogen or hydroxy,

          wherein

20           said phenyl and naphthyl have one or two substituents selected from the group consisting of fluoro, chloro, bromo, hydroxy, nitro, amino, N((C<sub>1-6</sub>) alkyl)amino, di(C<sub>1-6</sub>) alkylamino, N((C<sub>1-6</sub>) alkyl sulfonyl)-amino, morpholino, phenyl, pyridyl, trifluoromethyl, trifluoromethyloxy, (C<sub>1-6</sub>) alkoxy, and (C<sub>1-6</sub>) alkyl; and

25           R<sup>22</sup>   represents hydrogen or hydroxy.

30           (7)   The 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1 or 2,

wherein

R<sup>12</sup> and R<sup>13</sup> independently represent hydrogen, or (C<sub>1-6</sub>) alkyl optionally substituted by phenyl, naphthyl or pyridyl.

5

- (8) The 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1, wherein said derivative is selected from the group consisting of the following compounds:

10

N-[6-(benzyloxy)-2-naphthoyl]phenylalanine;

N-[6-(benzyloxy)-2-naphthoyl]-4-(trifluoromethyl)phenylalanine;

N-{6-[(4-fluorobenzyl)oxy]-2-naphthoyl}phenylalanine;

N-{6-[(3-fluorobenzyl)oxy]-2-naphthoyl}phenylalanine;

N-{6-[(2-fluorobenzyl)oxy]-2-naphthoyl}phenylalanine;

15

N-[6-(3-pyridinylmethoxy)-2-naphthoyl]phenylalanine;

N-{6-[(3,4-difluorobenzyl)oxy]-2-naphthoyl}phenylalanine;

N-{6-[2-(1H-pyrrol-1-yl)ethoxy]-2-naphthoyl}phenylalanine;

N-[6-(4-pyridinylmethoxy)-2-naphthoyl]phenylalanine;

N-[6-(benzyloxy)-2-naphthoyl]-3-(trifluoromethyl)phenylalanine;

20

N-[6-(benzyloxy)-2-naphthoyl]tryptophan;

N-[6-(benzyloxy)-2-naphthoyl]-O-methyltyrosine;

N-[6-(benzyloxy)-2-naphthoyl]-3-methoxytyrosine;

N-[6-(benzyloxy)-2-naphthoyl]-β-hydroxyphenylalanine;

N-[6-(2-phenylethoxy)-2-naphthoyl]phenylalanine;

25

N-[6-(benzyloxy)-2-naphthoyl]-4-chlorophenylalanine;

N-[6-(benzyloxy)-2-naphthoyl]-3-fluorophenylalanine;

N-{6-[(2-chlorobenzyl)oxy]-2-naphthoyl}phenylalanine;

N-{6-[(3-chlorobenzyl)oxy]-2-naphthoyl}phenylalanine;

N-{6-[(2-methoxybenzyl)oxy]-2-naphthoyl}phenylalanine;

30

N-{6-[(3-methoxybenzyl)oxy]-2-naphthoyl}phenylalanine;

N-{6-[(2,3-dichlorobenzyl)oxy]-2-naphthoyl}phenylalanine;

- 5 N-{6-[(3,5-dichlorobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3,5-dimethoxybenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(2-thienyl)alanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-bromophenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-nitrophenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-hydroxyphenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(1-naphthyl)alanine;  
N-[6-(benzyloxy)-2-naphthoyl]-5-hydroxytryptophan;  
N-[6-(benzyloxy)-2-naphthoyl]-2-fluorophenylalanine;  
10 N-{6-[(2-bromobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-bromobenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(2-methylbenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-methylbenzyl)oxy]-2-naphthoyl}phenylalanine;  
N-{6-[(3-nitrobenzyl)oxy]-2-naphthoyl}phenylalanine;  
15 N-[6-(benzyloxy)-2-naphthoyl]-3-(2-naphthyl)alanine;  
N-[6-(benzyloxy)-2-naphthoyl]-4-iodophenylalanine;  
N-[6-(benzyloxy)-2-naphthoyl]-5-fluorotryptophan;  
N-[6-(benzyloxy)-2-naphthoyl]-3-(1H-pyrrolo[2,3-b]pyridin-3-yl)alanine;  
N-{6-[2-(4-pyridinyl)ethoxy]-2-naphthoyl}phenylalanine;  
20 N-{6-[(3-ethoxybenzyl)oxy]-2-naphthoyl}phenylalanine; and  
N-[6-(2-phenylpropoxy)-2-naphthoyl]phenylalanine;

- 25 (9) A medicament comprising the 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 as an active ingredient.
- (10) The medicament as claimed in claim 9, further comprising one or more pharmaceutically acceptable excipients.

- (11) The medicament as claimed in claim 9, wherein the 2-naphthamide derivative, its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof is an IP receptor antagonist.
- 5 (12) The medicament as claimed in claim 9 for prophylaxis and/or treatment of urological disorder or disease.
- (13) The medicament as claimed in claim 9 for prophylaxis and/or treatment of pain.
- 10 (14) The medicament as claimed in claim 9 for prophylaxis and/or treatment of hypotension.
- (15) The medicament as claimed in claim 9 for prophylaxis and/or treatment of hemophilia and hemorrhage.
- 15 (16) The medicament as claimed in claim 9 for prophylaxis and/or treatment of inflammation.
- (17) Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prophylaxis of urological disorders.
- 20 (18) Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prophylaxis of pain.
- 25 (19) Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prophylaxis of hypotension.
- (20) Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prophylaxis of hemophilia and hemorrhage.
- 30

- (21) Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prophylaxis of inflammation.
- 5 (22) Process for controlling urological disorders in humans and animals by administration of an IP receptor-antagonistically effective amount of at least one compound according to claim 1.
- 10 (23) Process for controlling pain in humans and animals by administration of an IP receptor-antagonistically effective amount of at least one compound according to claim 1.
- 15 (24) Process for controlling hypotension in humans and animals by administration of an IP receptor-antagonistically effective amount of at least one compound according to claim 1.
- (25) Process for controlling hemophilia and hemorrhage in humans and animals by administration of an IP receptor-antagonistically effective amount of at least one compound according to claim 1.
- 20 (26) Process for controlling inflammation in humans and animals by administration of an IP receptor-antagonistically effective amount of at least one compound according to claim 1.

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP 03/05705

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C235/34 C07D209/20 C07D213/55 C07D333/24 A61K31/197  
A61K31/405 A61P13/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 68591 A (HOFFMANN LA ROCHE) 20 September 2001 (2001-09-20) page 2, line 3 -page 3, line 10 examples 1,2,6,7,13-15	1-26
A	WO 00 31045 A (KAWAMURA HIROYUKI ;TORII PHARMACEUTICAL CO LTD (JP); UCHIYAMA HIRO) 2 June 2000 (2000-06-02) cited in the application abstract page 84; example 79	1-10
A	WO 01 10383 A (BRENCHLEY GUY ;GOLEC JULIAN (GB); VERTEX PHARMA (US); CHARIFSON PA) 15 February 2001 (2001-02-15) examples 22,23,38	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

15 October 2003

Date of mailing of the international search report

28/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

O'Sullivan, P

## INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP 03/05705

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 719 760 A (ONO PHARMACEUTICAL CO) 3 July 1996 (1996-07-03) page 2, line 5 - line 18 tables -----	1-10



# INTERNATIONAL SEARCH REPORT

national application No. PCT/EP 03/05705

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 22-26 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 03/05705

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0168591	A	20-09-2001	AU 5216801 A	24-09-2001
			BR 0109235 A	17-12-2002
			CA 2401502 A1	20-09-2001
			CN 1418187 T	14-05-2003
			CZ 20023422 A3	15-01-2003
			WO 0168591 A1	20-09-2001
			EP 1265853 A1	18-12-2002
			HU 0300188 A2	28-06-2003
			JP 2003527368 T	16-09-2003
			NO 20024387 A	21-10-2002
			US 2001056100 A1	27-12-2001
WO 0031045	A	02-06-2000	AU 4058399 A	13-06-2000
			WO 0031045 A1	02-06-2000
WO 0110383	A	15-02-2001	AU 6894800 A	05-03-2001
			EP 1163208 A2	19-12-2001
			JP 2003506389 T	18-02-2003
			WO 0110383 A2	15-02-2001
			US 2002132833 A1	19-09-2002
EP 0719760	A	03-07-1996	AT 184871 T	15-10-1999
			DE 69512364 D1	28-10-1999
			DE 69512364 T2	03-02-2000
			DK 719760 T3	20-12-1999
			EP 0719760 A1	03-07-1996
			ES 2140629 T3	01-03-2000
			GR 3031808 T3	29-02-2000
			JP 8239356 A	17-09-1996
			US 6013673 A	11-01-2000
			US 5753700 A	19-05-1998

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**This Page Blank (uspto)**